

**PROPOSED**

**Species at Risk Act**  
Recovery Strategy Series

Recovery Strategy for Nooksack Dace  
(*Rhinichthys cataractae*) in Canada

# Nooksack Dace



Photo by: Mike Pearson

September 2006



Fisheries and Oceans  
Canada

Pêches et Océans  
Canada

Canada

## About the Species at Risk Act Recovery Strategy Series

### What is the *Species at Risk Act* (SARA)?

SARA is the Act developed by the federal government as a key contribution to the common national effort to protect and conserve species at risk in Canada. SARA came into force in 2003 and one of its purposes is *“to provide for the recovery of wildlife species that are extirpated, endangered or threatened as a result of human activity.”*

### What is recovery?

In the context of species at risk conservation, **recovery** is the process by which the decline of an endangered, threatened or extirpated species is arrested or reversed, and threats are removed or reduced to improve the likelihood of the species' persistence in the wild. A species will be considered **recovered** when its long-term persistence in the wild has been secured.

### What is a recovery strategy?

A recovery strategy is a planning document that identifies what needs to be done to arrest or reverse the decline of a species. It sets goals and objectives and identifies the main areas of activities to be undertaken. Detailed planning is done at the action plan stage.

Recovery strategy development is a commitment of all provinces and territories and of three federal agencies — Environment Canada, Parks Canada Agency and Fisheries and Oceans Canada — under the Accord for the Protection of Species at Risk. Sections 37–46 of SARA ([http://www.sararegistry.gc.ca/the\\_act/default\\_e.cfm](http://www.sararegistry.gc.ca/the_act/default_e.cfm)) spell out both the required content and the process for developing recovery strategies published in this series.

Depending on the status of the species and when it was assessed, a recovery strategy has to be developed within one to two years after the species is added to the List of Wildlife Species at Risk. Three to four years is allowed for those species that were automatically listed when SARA came into force.

### What's next?

In most cases, one or more action plans will be developed to define and guide implementation of the recovery strategy. Nevertheless, directions set in the recovery strategy are sufficient to begin involving communities, land users, and conservationists in recovery implementation. Cost-effective measures to prevent the reduction or loss of the species should not be postponed for lack of full scientific certainty.

### The series

This series presents the recovery strategies prepared or adopted by the federal government under SARA. New documents will be added regularly as species get listed and as strategies are updated.

### To learn more

To learn more about the Species at Risk Act and recovery initiatives, please consult the SARA Public Registry (<http://www.sararegistry.gc.ca/>) and the web site of the Recovery Secretariat ([http://www.speciesatrisk.gc.ca/recovery/default\\_e.cfm](http://www.speciesatrisk.gc.ca/recovery/default_e.cfm)).

**Recovery Strategy for Nooksack Dace  
(*Rhinichthys cataractae*) in Canada  
[Proposed]**

**September 2006**

**Recommended citation:**

Pearson, M.P., T. Hatfield, J.D. McPhail, J.S. Richardson, J.S. Rosenfeld, H. Schreier, D. Schluter, D.J. Snee, M. Stejovic, E.B. Taylor, and P.M. Wood. 2006. Recovery Strategy for Nooksack Dace (*Rhinichthys cataractae*) in Canada [Proposed]. Species at Risk Act Recovery Strategy Series. Vancouver: Fisheries and Oceans Canada. ix + 31pp.

**Additional copies:**

You can download additional copies from the SARA Public Registry (<http://www.sararegistry.gc.ca/>)

**Cover illustration:** M. Pearson

Egalement disponible en français sous le titre  
« Programme de rétablissement pour le naseux de Nooksack (*Rhinichthys cataractae*) au Canada [Projet] »

© Her Majesty the Queen in Right of Canada, represented by the Minister of Fisheries and Oceans, 2006.  
All rights reserved.

ISBN To come

Cat. no. To come

*Content (excluding the cover illustration) may be used without permission, with appropriate credit to the source.*

## **DECLARATION**

This proposed recovery strategy for Nooksack dace has been prepared in cooperation with the jurisdictions described in the Preface. Fisheries and Oceans Canada has reviewed and accepts this document as its recovery strategy for Nooksack dace as required by the *Species at Risk Act*.

Success in the recovery of this species depends on the commitment and cooperation of many different constituencies that will be involved in implementing the directions set out in this strategy and will not be achieved by Fisheries and Oceans Canada or any other jurisdiction alone. In the spirit of the National Accord for the Protection of Species at Risk, the Minister of Fisheries and Oceans invites all Canadians to join Fisheries and Oceans Canada in supporting and implementing this strategy for the benefit of Nooksack dace and Canadian society as a whole. Fisheries and Oceans Canada will support implementation of this strategy to the extent possible, given available resources and its overall responsibility for species at risk conservation. The Minister will report on progress within five years.

This strategy will be complemented by one or more action plans that will provide details on specific recovery measures to be taken to support conservation of the species. The Minister will take steps to ensure that, to the extent possible, Canadians interested in or affected by these measures will be consulted.

## **RESPONSIBLE JURISDICTIONS**

The responsible jurisdiction for Nooksack dace under the *Species at Risk Act* is Fisheries and Oceans Canada. The Province of British Columbia also collaborated in the development of this recovery strategy.

## **AUTHORS**

DFO and the Province of British Columbia cooperated in the development of this recovery strategy. A recovery team was assembled to provide science-based recommendations to government with respect to the recovery of Nooksack dace. Members of the National Recovery Team for Nooksack Dace are listed below:

Todd Hatfield, Solander Ecological Research (Coordinator)  
Don McPhail, University of British Columbia  
Mike Pearson, Pearson Ecological (Writer)  
John Richardson, University of British Columbia  
Jordan Rosenfeld, British Columbia Ministry of Environment (Co-Chair)  
Hans Schreier, University of British Columbia  
Dolph Schluter, University of British Columbia  
Dan Sneep, Fisheries and Oceans Canada (Co-Chair)

Marina Stejpovic, Township of Langley  
Eric Taylor, University of British Columbia  
Paul Wood, University of British Columbia

## **ACKNOWLEDGMENTS**

Fisheries and Oceans Canada and the Province of BC are grateful to the technical experts involved in drafting this strategy, for their time and effort in attending meetings and reviewing the document. Financial support for the development of the recovery strategy was provided by the Habitat Conservation Trust Fund and the Province of British Columbia.

## **STRATEGIC ENVIRONMENTAL ASSESSMENT**

A strategic environmental assessment (SEA) is conducted on all SARA recovery planning documents, in accordance with the *Cabinet Directive on the Environmental Assessment of Policy, Plan and Program Proposals*. The purpose of a SEA is to incorporate environmental considerations into the development of public policies, plans, and program proposals to support environmentally-sound decision making.

Recovery planning is intended to benefit species at risk and biodiversity in general. However, it is recognized that strategies may also inadvertently lead to environmental effects beyond the intended benefits. The recovery planning process based on national guidelines directly incorporates consideration of all environmental effects, with a particular focus on possible impacts on non-target species or habitats. The results of the SEA are incorporated directly in the strategy itself, but are also summarized below.

While this recovery strategy will clearly benefit the environment by promoting the recovery of Nooksack dace, some potentially adverse effects on other species were also considered. The strategy calls for the protection, creation, and enhancement of riffle habitat, which could require control of beavers and their dams, and which might eliminate some of the deep pool and marsh habitat of Salish sucker, another species listed as Endangered under SARA. The strategy recommends cooperation with local stewardship groups and agency staff on beaver management, and proposes to address potential conflicts with recovery of Salish sucker by coordinating recovery activities for both species in watersheds where they coexist through the development of a joint Action Plan. The recovery strategy also calls for minimization of impacts of introduced predators, through documenting their occurrence and educating the public on their impacts. Further information on potential interactions with other species is presented in the Recovery section of the document, in particular under the headings *Broad Strategies to Reduce Threats and Effects on Other Species*. Taking these approaches into account, it was concluded that the benefits of this recovery strategy far outweigh any adverse effects that may result.

## RESIDENCE

SARA defines residence as: “a dwelling -place, such as a den, nest or other similar area or place, that is occupied or habitually occupied by one or more individuals during all or part of their life cycles, including breeding, rearing, staging, wintering, feeding or hibernating” [SARA S2(1)].

Residence identification is part of the listing process. Residence descriptions, or the rationale for why the residence concept does not apply to a given species, are posted on the SARA public registry:

[http://www.sararegistry.gc.ca/plans/residence\\_e.cfm](http://www.sararegistry.gc.ca/plans/residence_e.cfm)

## PREFACE

The Nooksack dace is a freshwater fish, under the jurisdiction of the federal government. The *Species at Risk Act* (SARA, Section 37) requires the competent minister to prepare recovery strategies for listed Extirpated, Endangered or Threatened species. The Nooksack dace was listed as Endangered under SARA in June 2003. Fisheries and Oceans Canada - Pacific Region and the Province of British Columbia co-led the development of this recovery strategy. The proposed strategy meets SARA requirements in terms of content and process (Sections 39-41). It was developed in cooperation or consultation with:

- The University of British Columbia
- The Township Of Langley

## **EXECUTIVE SUMMARY**

### **Background**

The Nooksack dace is a small (<15 cm) stream-dwelling cyprinid (minnow). It is considered a subspecies of the widespread and common longnose dace *Rhinichthys cataractae*. Within Canada it is known from four lowland streams in British Columbia's Fraser Valley. The global distribution includes approximately 20 additional streams in north-west Washington (McPhail 1997). The Nooksack dace is extirpated from some tributaries in Canadian watersheds where it was abundant in the 1960s (McPhail 1997). Its current status in Washington State is unknown.

Nooksack dace are strongly associated with riffle habitats (McPhail 1997) and the proportion of riffle in a reach is the strongest predictor of their presence (Pearson 2004a). Young-of-the-year fish require shallow pool habitats in close proximity to the riffles inhabited by adults (McPhail 1997). Home range size is typically very small (<50 m of channel) although a few individuals venture for at least hundreds of metres (Pearson 2004a). This suggests that clusters of riffles may contain semi-isolated subpopulations and that metapopulation dynamics may be important at the watershed scale (Pearson 2004a).

### **Threats**

Nooksack dace populations appear to be most vulnerable to seasonal lack of water, habitat loss to drainage activities, sediment deposition, and riffle loss to beaver ponds. Introduced predators are widespread in the range but probably have minimal impacts on Nooksack dace because of lack of habitat overlap. Hypoxia and toxicity are significant threats in some sections of at least one watershed, but do not threaten the species throughout its range.

### **Critical Habitat**

Critical habitat for Nooksack dace has not been formally described in this recovery strategy. The Recovery Team has compiled scientific data which will assist in the definition of critical habitat, and this information should provide the basis for an official designation of critical habitat through the action planning process, which will include socioeconomic analysis and consultation with affected interests. Critical habitat for Nooksack dace should be defined at the reach scale, and should include specific features such as riffles, shallow pools, and riparian habitat. Further studies are required to confirm the presence of other Nooksack dace populations and their critical habitats, and to characterize specific threats. Defining critical habitat will contribute to the refinement of recovery objectives and the management of activities that impact the species.

### **Recovery**

Recovery of Nooksack dace populations is both technically and biologically feasible. It will involve the establishment and/or maintenance of sufficient high quality riffle habitat in each creek to maintain a population. Specific requirements will vary, but will generally include in-stream flow protection, restoration of riffle habitat and, in



some circumstances, restriction of beaver impoundment. Some management will be required in all watersheds.

The goal of recovery is:

*To ensure long-term viability of Nooksack dace populations throughout their natural distribution in Canada.*

The recovery strategy has three objectives, each of which is discussed in detail in the text.

1. For all *currently and historically suitable* habitats in native streams to be occupied by 2015.
2. To increase Nooksack dace abundance to target levels in all watersheds by 2015.
3. To ensure that at least one reach in each watershed supports a high density of Nooksack dace.

Eight broad strategies have been identified in support of these objectives:

1. Protect, create and enhance riffle habitat in habitat reaches with high potential productivity.
2. Establish or maintain adequate baseflow in all habitats with high potential productivity.
3. Reduce sediment entry to creeks.
4. Ensure the integrity and proper functioning of riparian zones throughout watersheds.
5. Reduce habitat fragmentation.
6. Encourage stewardship amongst private landowners and the general public.
7. Minimize toxic contamination of creeks.
8. Minimize impacts of introduced predators.

# TABLE OF CONTENTS

EXECUTIVE SUMMARY .....	VIII
BACKGROUND .....	1
SPECIES INFORMATION .....	1
SPECIES DESCRIPTION .....	1
POPULATIONS AND DISTRIBUTION.....	1
DESCRIPTION OF THE SPECIES NEEDS.....	2
THREATS .....	5
IDENTIFICATION OF THE THREATS TO THE SURVIVAL OF THE SPECIES .....	5
Threat 1: Physical Destruction of Habitat.....	6
Threat 2: Seasonal Lack of Water .....	8
Threat 3: Sediment Deposition .....	8
Threat 4: Riffle Loss to Beaver Ponds .....	9
Threat 5: Habitat Fragmentation .....	9
Threat 6: Toxicity .....	10
Threat 7: Hypoxia .....	11
Threat 8: Increased Predation .....	11
SUMMARY OF THREATS ANALYSIS .....	12
CRITICAL HABITAT .....	12
IDENTIFICATION OF CRITICAL HABITAT .....	12
SCHEDULE OF STUDIES TO IDENTIFY CRITICAL HABITAT .....	14
KNOWLEDGE GAPS IN NOOKSACK DACE BIOLOGY .....	14
RECOVERY .....	15
RECOVERY FEASIBILITY .....	15
RECOVERY GOAL, OBJECTIVES AND CORRESPONDING APPROACHES .....	16
Recovery Goal:.....	16
Recovery Objectives.....	16
Broad Strategies to be Taken to Address Threats .....	17
Evaluation .....	18
Effects on Other Species .....	18
APPROACHES TO RECOVERY .....	18
ACTIONS ALREADY COMPLETE OR UNDERWAY .....	19
STATEMENT OF WHEN AN ACTION PLAN WILL BE COMPLETED.....	26
LITERATURE CITED.....	27
APPENDIX: RECORD OF COOPERATION AND CONSULTATION.....	31

## BACKGROUND

### SPECIES INFORMATION

The status report and assessment summary for Nooksack Dace is available from the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) Secretariat ([www.cosewic.gc.ca](http://www.cosewic.gc.ca)).

<b>Common Name:</b>	Nooksack Dace
<b>Scientific Name:</b>	<i>Rhinichthys cataractae</i>
<b>Assessment Summary:</b>	May 2000
<b>COSEWIC Status:</b>	Endangered, April 1996
<b>SARA Status:</b>	Endangered, June 2003
<b>Reason for Designation:</b>	This species has a restricted range in Canada, and is in significant decline due to habitat loss and degradation.
<b>Range in Canada:</b>	British Columbia
<b>Status History:</b>	Designated Endangered in April 1996. Status re-examined and confirmed in May 2000. Last assessment based on an existing status report.

#### Species Description

The Nooksack dace is a small (<15 cm) stream dwelling cyprinid (minnow). The body is streamlined with, large pectoral fins and a snout that overhangs the mouth. Body colouration is grey-green above a dull, brassy lateral stripe and dirty white below it. There is often a distinct black stripe on the head in front of the eyes, which in juveniles continues down the flanks to the tail (McPhail 1997). The Nooksack dace is considered a subspecies of the widespread and common longnose dace *Rhinichthys cataractae* (J.D. McPhail, University of British Columbia, pers. comm.). It evolved through geographic isolation in Washington State's Chehalis River valley sometime during the Pleistocene glaciations (McPhail 1997). Adults are generalized insectivores while juveniles feed on zooplankton (McPhail 1997).

#### Populations and Distribution

Populations are documented from four lowland streams in British Columbia's Fraser Valley (Figure 1). The global distribution consists of approximately 20 additional streams in north-west Washington State. The species is extirpated from some tributaries within Canadian watersheds where it was abundant in the 1960s (McPhail 1997). The current status of Washington State populations is unknown. Based on

available information, Canada contains approximately 10% of the global range and 20% of all populations (Figure 1).

### **Description of the Species Needs**

#### **Biological Needs, Ecological Role and Limiting Factors**

The major factor limiting population abundance and distribution is the availability of high quality habitat (see below). Given adequate habitat Nooksack dace populations should recover rapidly as their life history characteristics promote rapid population growth. They are small-bodied, mature early (2 years, McPhail 1997), and have an extended spawning period and may spawn more than once each year (April - July, Pearson 2004a), a trait that increases fecundity in species otherwise limited by small female body size (Blueweiss et al. 1978; Burt et al. 1988).




Figure 1: Canadian and global distribution of Nooksack dace. In Canada the Nooksack dace is known to inhabit four watersheds (left panel; 1- Brunette River, 2 – Bertrand Creek, 3 – Pepin Brook, 4 – Fishtrap Creek). Globally, it is also found in a number of other streams in northwestern Washington (right panel, adapted from McPhail 1997).

## Habitat Needs

### ***Physical Habitat***

Nooksack dace are riffle specialists. The proportion of riffle habitat in a reach is the strongest predictor of their presence and they are rarely found in reaches with less than 10 percent riffle by length (Figure 2) or in reaches where long stretches of deep pool habitat separate riffles (Pearson 2004a). Young-of-the-year fish require shallow calm, pool habitats in close proximity to riffles. Most individuals appear to have small home ranges (tens of metres of channel) although a small number of individuals venture hundreds of metres. Clusters of riffles may contain semi-isolated subpopulations. Distances and barriers between clusters may influence long-term population persistence by altering watershed scale population dynamics.




Figure 2: Nooksack dace are found in fewer than half of reaches that contain less than 10 percent riffle by length. Numbers over bars indicate sample size (adapted from Pearson 2004a).

### ***Water Quantity***

Riffles are among the shallowest of stream habitats and consequently among the first to shrink when flow declines. When surface flow ceases, riffle habitat is entirely eliminated and Nooksack dace may be forced into pools, a non-preferred habitat where foraging success and security from predation may be compromised.

### ***Water Quality***

Little information exists on tolerances or preferences of Nooksack dace for parameters such as dissolved oxygen, pH, and temperature. Activity appears minimal at temperatures below 11 °C, and fish forage normally at temperatures in excess of 20 °C (Pearson 2004a). Nooksack dace are likely poorly adapted to

hypoxia as their riffle habitats are typically well oxygenated. The federal water quality guideline for dissolved oxygen to support aquatic life (5 mg/l, CCREM 1987) is an appropriate benchmark for habitat assessment.

## THREATS

### Identification of the threats to the survival of the species

The prospects for recovery of a species at risk depend upon its vulnerability to the threats facing it, their severity and ubiquity across the range. In the following sections we summarize detailed analyses of each of these factors that are published elsewhere (Pearson 2004a, 2004b).

Eight factors (Table 1) are considered threats based on knowledge of species biology and habitat conditions across the Canadian range. All are proximate, in that they act directly upon the fish or their habitats. The vulnerability of Nooksack dace to each threat, and the severity of each threat in each watershed are rated and summarized graphically in Table 2. The ratings are based on analyses of a suite of factors that cause, exacerbate, or mitigate threats (Figure 3), and are briefly summarized in the text. A summary by watershed is presented in Table 3. For details of assessment methods and rationales for ratings see Pearson (2004a, 2004b).

Table 1: Potential threats to Nooksack dace in Canada in descending order of concern.

Threat	Management Concern
1. Physical Destruction of Habitat:	Drainage, dyking, channelization and infilling of water bodies destroying habitat.
2. Seasonal Lack of Water:	Low flows in late summer eliminate habitat, reducing fitness or survival.
3. Sediment Deposition:	Deposited sediment degrading habitat.
4. Riffle Loss to Beaver Ponds:	Beaver ponds flooding riffle habitat.
5. Habitat Fragmentation:	Permanent or temporary barriers preventing or inhibiting fish from traversing some stream reaches. This restricts access to usable habitats and/or alters metapopulation dynamics to increase extinction risk.
6. Toxicity:	Toxic discharges from point or non-point sources significantly reducing survival or fitness.
7. Hypoxia:	Episodes of extreme hypoxia causing acute mortality or reduced fitness.
8. Increased Predation:	Introduced predators consuming individuals or reducing their fitness by inducing behavioural changes.

Table 2: Summary of threats assessment for Nooksack dace (see text for basis of assessment).

<i>Threat</i>	Vulnerability of Nooksack Dace	Severity Across Range
Physical Destruction of Habitat	***	***
Seasonal Lack of Water	***	***
Sediment Deposition	***	***
Riffle Loss to Beaver Ponds	***	**
Habitat Fragmentation	**	***
Toxicity	**	**
Hypoxia	*	**
Increased Predation	*	***

***	major concern	**	moderate concern	*	minor concern
-----	---------------	----	------------------	---	---------------

Table 3: Assessment of threat severity in each of the four watersheds from which Nooksack dace are known in Canada. Background data and details of assessment methods for Bertrand, Pepin and Fishtrap Creeks are provided by Pearson (2004a). The Brunette River population was discovered in 2004 and a threats analysis has not been completed for it.

<i>Threat</i>	Bertrand Creek	Pepin Brook	Fishtrap Creek	Brunette River
Hypoxia	**	***	**	?
Physical Destruction of Habitat	**	***	***	?
Habitat Fragmentation	***	**	**	?
Toxicity	**	*	***	?
Sediment Deposition	**	***	**	?
Seasonal Lack of water	***	*	**	?
Increased Predation	**	**	**	?
Riffle Loss to Beaver Ponds	*	***	*	?

### ***Threat 1: Physical Destruction of Habitat***

#### *Description*

Channelization, dredging and infilling directly destroying or degrading stream habitats.

#### *Vulnerability (major concern)*

The riffle habitats required by Nooksack dace are the ‘high spots’ in a stream, and tend to be targeted for removal or alteration in drainage projects. Channelization and drainage work also typically eliminates the shallow marginal pools preferred by young-of-the-year.

#### *Severity (major concern)*



Approximately 77% of pre-settlement wetland areas in the Fraser Valley have been drained or infilled (Boyle et al. 1997). Fifteen percent of the area's streams no longer exist, having been paved over or piped (Fisheries and Oceans Canada 1998). A large, but unknown, proportion of those that remain have been channelized and/or repeatedly dredged in agricultural drainage or urban development projects. It is




Figure 3: Factors known or suspected to drive or trigger threats to Nooksack dace (from Pearson 2004b).

difficult to overstate the historical extent of fish habitat loss to these activities. Both permitted and un-permitted dredging of ditches and stream channels for flood control and agricultural drainage still occur annually in all watersheds included in this strategy. In recent years, Fishtrap Creek has been most affected. The lower 5 km of the mainstem were dredged by the City of Abbotsford in 1990-1991 (Pearson 2004a), eliminating riffle from what was previously a densely populated reach (J.D. McPhail, UBC pers. comm.).

### ***Threat 2: Seasonal Lack of Water***

#### *Description*

During late summer, when rainfall is sparse, Fraser Valley stream flows are maintained almost solely by groundwater. Stream hydrographs vary widely depending on surface soil permeability and water use. Watersheds with large unconfined aquifers maintain steady flows of cold water throughout this critical period, while surface flows may cease completely in watersheds with impermeable surface soils. Unfortunately the late summer low-flow period coincides with peak demand for water withdrawal from wells and streams for irrigation and domestic use. Common land use changes in the Fraser Valley also tend to exacerbate problems with water availability. Gravel mining reduces the size of the aquifer contributing to baseflow, urban development increases the area of impermeable surfaces (reducing infiltration to the aquifer), and agricultural drainage lowers water tables, further reducing flows.

#### *Vulnerability (major concern)*

Nooksack dace are highly vulnerable to lack of water. Adults inhabit riffles and young-of-the-year school in nearby shallow pools (McPhail 1997). These habitats are the first to be affected by lack of water. Adults also spawn in riffles, but during spring and early summer when water is more plentiful.

#### *Severity (major concern)*

Low surface flows have reduced the availability of suitable habitat in Bertrand and Fishtrap creeks for several weeks during very dry years (Pearson, pers. obs.). Nooksack dace are especially vulnerable to further wetland drainage, increases in impermeable surfaces and/or water withdrawal. Extensive gravel mining is underway in two watersheds and will reduce baseflow in these systems by an unknown amount in future.

### ***Threat 3: Sediment Deposition***

#### *Description*

Sediment deposition is controlled by the balance between the rate of sediment delivery to the channel and capacity of the stream to mobilize and carry it downstream. Sediment delivery may be increased by direct discharges, storm drain runoff, or bank erosion accelerated by lack of riparian vegetation and/or increased peak flows (Waters 1995). All of these sources are likely to increase with urban, agricultural and mining development in a watershed.

*Vulnerability (major concern)*

Adult dace spawn, forage and rest in the crevasses between and under coarse riffle substrate (McPhail 1997). Sedimentation clogs these spaces and inhibits the flow of oxygenated water through the substrate. It is less likely to be a problem for young-of-the-year dace, which inhabit the water column in shallow pools (McPhail 1997).

*Severity (major concern)*

Significant sediment deposition occurs in portions of all watersheds (Pearson 2004a).

***Threat 4: Riffle Loss to Beaver Ponds***

*Description*

Beaver ponds have been shown to influence fish populations both positively and negatively (Hanson & Campbell 1963; Keast & Fox 1990; Lavkulich et al. 1999; Schlosser 1995). The impacts of riffle loss through ponding have received scant attention, but may be significant for species like Nooksack dace, which depend on these habitats.

*Vulnerability (major concern)*

Nooksack dace are riffle specialists. The proportion of riffle habitat a reach contains is the best predictor of their presence, and dace are absent from long sections of continuous deep pool, like beaver ponds, even when riffles are present (Pearson 2004a).

*Severity (moderate concern)*

Riffle loss to beaver ponding is a major concern in at least one watershed, Pepin Brook. In 1999, beavers had impounded 47% of its 6.4 km mainstem. By 2001 an additional 690 m of channel was impounded, eliminating 10% of the 938 m of riffle recorded in the 1999 survey (Pearson 2004a). Impounded area did not change in two other watersheds monitored over the same period (Bertrand and Fishtrap creeks) as higher winter flows washed out dams regularly and narrower riparian forest strips probably limit the food supply of beavers (Pearson 2004a).

***Threat 5: Habitat Fragmentation***

*Description*

Physical barriers such as perched culverts, beaver dams, and agricultural weirs commonly prevent movement between habitats for all or part of the year in Fraser Valley streams. In addition, any of the other threats discussed may fragment habitat by preventing or curtailing movement of fish through affected reaches. On a larger scale, connections between watersheds during floods were undoubtedly more common prior to the extensive dyking and drainage works of the past century.

*Vulnerability (moderate concern)*

Most Nooksack dace have very small home ranges, covering less than 50 m of channel, although a few individuals appear to venture further (Pearson 2004a). The distribution of populations is also very clumped within streams. In combination, these

data suggest that each watershed is inhabited by loosely connected subpopulations. Most barriers and habitat fragmentation in Nooksack dace watersheds date from 50 to 130 years ago, and surviving populations have shown some resilience (Pearson 2004a). The effects of less movement between subpopulations and reduced ability to colonize new habitat, however, may occur over longer time frames. The extent and importance of this to the long-term persistence of individual subpopulations and to recolonization following local extinctions of subpopulations is unclear.

*Severity (major concern)*

The extensive destruction of aquatic habitat that has occurred within the Fraser Valley over the past 150 years (see *Physical Destruction of Habitat* above) has fragmented habitat badly. Within watersheds, physical barriers and degraded habitat have likely affected movement patterns between subpopulations. Bertrand, Pepin and Fishtrap Creeks are all tributaries of the Nooksack River, but are isolated from one another by poor habitat conditions in the Washington State portion of their watersheds (McPhail 1997). Fish and habitat distributions within the Brunette system have yet to be surveyed.

***Threat 6: Toxicity***

*Description*

Toxic compounds enter Fraser Valley streams through urban storm runoff, contaminated groundwater (e.g. agricultural pesticides and herbicides), direct industrial discharges, sewage treatment plant effluents, aerial deposition, and accidental spills (Hall et al. 1991). Concentrations in the water column are widely variable over time because dilution varies with stream discharge and inputs are often pulsed (e.g. first flush of stormwater following a long dry spell, Hall et al. 1991). Some contaminants, particularly heavy metals, bind to sediments where they may be taken up and bioaccumulated by aquatic invertebrates and subsequently fish.

*Vulnerability (moderate concern)*

Data on threshold concentrations for lethal and sublethal effects of toxic compounds on Nooksack dace are lacking. As a bottom-dwelling species, they may be sensitive to contaminants bound to sediment as well as those in food items and the water column.

*Severity (moderate concern)*

Toxicity is likely to impact some Nooksack dace populations. Large portions of the Fishtrap Creek, Bertrand Creek, and the Brunette River watersheds are urbanized, which generally causes elevated levels of copper, lead and zinc in stream sediments (Hall et al. 1991). Row crop agriculture with intensive pesticide/herbicide use is also common in the Fishtrap Creek watershed (Pearson 2004a). The range of compounds that could enter creeks from spraying, poor waste management, and accidental spills is enormous.

### **Threat 7: Hypoxia**

#### *Description*

Hypoxia is ultimately caused by the cumulative effects of local and watershed-scale impacts. Increased nutrients result in algal blooms and high densities of macrophytes that strip the water of oxygen at night. Decomposition of dead algae and vegetation exacerbates the problem and may severely depress daytime oxygen levels as well. Nutrients in Fraser Valley groundwater and streams are elevated, primarily a consequence of over-application of manure and fertilizers to agriculture lands (Lavkulich et al. 1999; Schreier et al. 2003), but also of urban stormwater runoff and septic systems (Lavkulich et al. 1999). Lack of shade from riparian vegetation permits water temperatures to rise. Warmer water has less capacity for dissolved oxygen and increases the metabolic demands of fish and other organisms. Reduced water movement impairs reoxygenation of water and may be caused by channelization, (Schreier et al. 2003), beaver ponds (Fox & Keast 1990; Schlosser & Kallemyn 2000), or low flows.

#### *Vulnerability (minor concern)*

Lethally low levels of hypoxia are unknown for Nooksack dace, but riffles are generally well-oxygenated habitats and species that are specialized to inhabit them are unlikely to be well adapted to hypoxia. Even moderate levels of chronic hypoxia may reduce growth, condition, and fecundity. In the absence of better information, the federal guideline for the protection of aquatic life ( $5\text{mg}\cdot\text{l}^{-1}$ , CCREM 1987) is a useful target.

#### *Severity (moderate concern)*

Hypoxia is a major concern in at least one stream, Pepin Brook, and a moderate concern in Bertrand and Fishtrap Creeks. Fish inhabiting riffles and shallow pools immediately below hypoxic reaches may be affected, although this comprises a small proportion of total habitat.

### **Threat 8: Increased Predation**

#### *Description*

Increased predation is most likely to arise from the introduction of new species to Nooksack dace habitats. Such introductions are implicated in the extinction of numerous native fishes across North America (Gido & Brown 1999; Miller et al. 1989; Richter 1997).

#### *Vulnerability (minor concern)*

The impacts of introduced predators on Nooksack dace populations are unknown. Populations have coexisted with bullheads (*Ameiurus nebulosis*), bullfrogs (*Rana catesbeiana*), smallmouth bass (*Micropterus dolomieu*), or largemouth bass (*M. salmoides*) for at least ten years in these watersheds (Pearson, unpubl.). All of these species would undoubtedly prey upon Nooksack dace given the opportunity, but there is little habitat overlap. These predators thrive in warm water littoral zones (Corkran & Thoms 1996; Scott & Crossman 1973) and are very rarely found in riffles. Lack of water could, however, force Nooksack dace out of riffles and into

pools where predation risk is likely to be much higher. The possibility of a new, effective predator being introduced to Nooksack dace habitat is also ever present.

*Severity (major concern)*

Introduced predators inhabit every stream known to contain Nooksack dace.

**Summary of Threats Analysis**

Nooksack dace populations appear to be most vulnerable to seasonal lack of water, habitat loss to drainage activities, sediment deposition, and riffle loss to beaver ponds. Habitat fragmentation is likely having some impacts in all watersheds and is considered a moderate concern. Introduced predators are widespread in the range but probably have minimal impacts on Nooksack dace because of lack of habitat overlap. Hypoxia and toxicity are significant threats in some sections of at least one watershed, but do not threaten the species throughout its range.

**CRITICAL HABITAT**

**Identification of Critical Habitat**

Critical habitat is defined in SARA as “the habitat that is necessary for the survival or recovery of a listed wildlife species and that is identified as the species’ critical habitat in the recovery strategy or in an action plan for the species.” [SARA S. 2(1)]. Critical habitat for Nooksack dace has not been formally described in this recovery strategy, though the Recovery Team has compiled scientific data which will assist in the scientific definition of critical habitat, including a separate assessment of the habitat requirements of the species (National Recovery Team for Salish Sucker and Nooksack Dace 2005). Specific features of critical habitat for Nooksack dace are summarized below. This information should provide the basis for an official designation of critical habitat through the action planning process, which will include socioeconomic analysis and consultation with affected interests. Defining critical habitat will contribute to the refinement of recovery objectives and the management of activities that impact the species.

**Critical Habitat Features**

Based on available physical and biological data, Nooksack dace critical habitat could include the following key elements:

***The Reach Scale***

Riffles and shallow pools (see below) are the required habitats of Nooksack dace, but critical habitat should be defined at the reach scale, a larger, natural unit of river morphology that ranges from hundreds to thousands of metres in length (Frissell et al. 1986). There are three reasons for adopting this scale. First, the reach scale corresponds to the distribution of subpopulations within watersheds (Pearson 2004a). Second, the ‘channel units’ of critical habitat (riffles and shallow pools) are dynamic and frequently move during flood events in these streams. In Bertrand

Creek, this occurs on an annual basis (Pearson pers. obs.). Effective protection and management of critical habitat in these circumstances must allow for normal channel processes and must, therefore, occur at a spatial scale larger than the channel unit. The reach scale is the next largest in accepted stream habitat classifications (Frissell et al. 1986; Imhof et al. 1996). Third, the reach scale corresponds most closely to that of land ownership in these watersheds and, consequently, to most potential recovery actions.

### ***Riffle Habitat***

Available information overwhelmingly suggests that riffles are critical to species persistence. Nooksack dace typically occur in riffles over loose gravel and cobble substrates where water velocity exceeds  $0.25 \text{ m}\cdot\text{s}^{-1}$ . They spawn near the upstream end of riffles (McPhail 1997) between late April and early July (Pearson 2004a) and forage nocturnally for riffle dwelling insects (McPhail 1997). The percent of riffle in a stream reach is a good predictor of dace presence. Riffles that are isolated by long stretches of deep pool, however, are seldom inhabited (Pearson 2004a). A threshold of 10% riffle by length would exclude these small isolated riffles that have little value to Nooksack dace.

### ***Shallow Pool Habitat***

Young-of-the-year Nooksack dace inhabit shallow (10-20 cm) pools adjacent to riffles where they swim above sand, mud, or leaf litter substrates and feed upon chironomid pupae and ostracods (McPhail 1997). Loss of these habitats will quickly produce population-level impacts.

### ***Riparian Habitat***

Riparian vegetation should be included in critical habitat to the extent it is necessary to protect the integrity of in-stream critical habitat. Required widths would vary among sites and should be defined in reach scale assessments. Reserves must be sufficient to control sediment entry to the stream from overland flow, to prevent excessive bank erosion and to buffer stream temperatures. Reserve areas will also remove significant amounts of nitrate and phosphorous from groundwater, although their efficiency depends strongly on hydrogeologic conditions (Martin et al. 1999; Puckett 2004; Wigington et al. 2003). The effectiveness of a riparian reserve in preventing materials (e.g., sediments, nutrients, toxins) from entering a stream depends strongly on its continuity in addition to its width (Weller et al. 1998). Consequently, riparian reserves in critical habitat reaches should be continuous. In open landscapes, such as agricultural fields, vegetation from reserve areas will collect windblown insects (Whitaker et al. 2000). Such insects, falling from riparian vegetation into the water constitute an important food source in headwater streams (Allan et al. 2003; Schlosser 1991).

It is important to understand that in some circumstances, more than 30 m of riparian vegetation may be required for full mitigation of warming (Brown & Krygier 1970; Castelle et al. 1994; Lynch et al. 1984) and siltation (Davies & Nelson 1994; Kiffney et al. 2003; Moring 1982), and for long-term maintenance of channel morphology

(Murphy et al. 1986; Murphy & Koski 1989). At least 10 m are required to maintain levels of terrestrial food inputs similar to those of forested landscapes (Culp & Davies 1983). Reserves as narrow as 5 m provide significant protection from bank erosion and sediment deposition from overland flow (Lee et al. 2003; McKergow et al. 2003).

Failure to maintain an adequate riparian reserve as part of critical habitat would be highly likely to cause population-level impacts. In habitats lacking sufficient flow or groundwater sources, lack of shade may increase water temperatures to harmful levels. Increased erosion due to poorer bank stability will cause sediment deposition in riffles, impairing spawning and incubation, reducing food availability, and eliminating the interstitial spaces in coarse substrate that dace occupy. Nutrient loading will be higher in reaches without adequate riparian vegetation (Dhondt et al. 2002; Lee et al. 2003; Martin et al. 1999) and is likely to contribute to hypoxia through eutrophication. Solar radiation will also be higher in reaches lacking adequate riparian shading (Kiffney et al. 2003) and will contribute to eutrophication. Reserves of 30 m or more should be maintained around Nooksack dace habitat wherever feasible to provide a high level of protection from impacts of adjacent land uses.

### Schedule of Studies to Identify Critical Habitat

Information exists to assist in the definition of critical habitat for Nooksack dace throughout its presently known range. Further surveys are required to identify other potential populations and characterize their critical habitats, as summarized below:

Study	Description	Timeframe	Status
Population Identification	The Coquitlam and Alouette Rivers are suspected of containing Nooksack dace based on a preliminary genetic and morphometric study of their <i>R. cataractae</i> populations (J.D. McPhail, UBC, unpubl. data). Additional samples are required for confirmation.	2005-2006	Underway
Critical Habitat Surveys	Habitat in the Brunette River has not been surveyed as its populations were unknown prior to 2004. Surveys will also be required in the Coquitlam and Alouette Rivers if the presence of Nooksack dace is confirmed there.	2006-2007	Planned

### Knowledge Gaps in Nooksack Dace Biology

Additional studies should be conducted to address the following data needs related to specific threats to Nooksack dace. This information will contribute to the protection of Nooksack dace and their critical habitats.

Study	Description	Timeframe	Status
Impacts of Riffle Drying	The fate of dace in reaches that dewater during late summer is uncertain. Sampling during this period will resolve whether fish leave the reach, move into pools, burrow into substrate, or die.	2004-2005	Underway



Impacts of Sediment Deposition in Riffles	The extent to which sediment deposited in riffles affects their ability to support healthy dace populations is uncertain and needs to be quantified.	2007-2008	Need Identified
---	--	-----------	-----------------

## RECOVERY

### Recovery Feasibility *Feasibility Criteria*<sup>1</sup>.

1. *Are individuals capable of reproduction currently available to improve the population growth or population abundance?*  
Yes. Breeding adults have been captured recently from all populations.
2. *Is sufficient habitat available to support the species or could it be made available through habitat management or restoration?*  
Yes. Sufficient physical habitat exists to support the three populations that have been surveyed (Bertrand, Pepin and Fishtrap creeks), although up to 70% of it is seriously degraded by sediment deposits or low water levels in late summer. The severity and extent of these problems could be mitigated by reducing ground and surface water withdrawals during sensitive periods, by reducing sediment entry to streams and by managing beaver activity in sensitive habitats. The quantity and condition of available habitat in the Brunette River population is unknown at present.
3. *Can significant threats to the species or its habitats be avoided or mitigated through recovery actions?*  
Yes. Riffle degradation through seasonal drying can be avoided by reducing water withdrawals or flow supplementation. Sedimentation can be reduced through riparian planting, improved agricultural practices, the installation of sediment traps in storm sewer systems, and proper sediment control at mine and construction sites. Riffle loss can be mitigated through habitat restoration and (when necessary) beaver control.
4. *Do the necessary recovery techniques exist and are they demonstrated to be effective?*  
Yes. Techniques to reduce problems of low base flow, sediment deposition and beaver ponding are well known. Monitoring of created riffle habitat has demonstrated that restored habitats are quickly colonized.

### **Feasibility Assessment**

Recovery of Nooksack dace populations to levels ensuring long-term survival is both technically and biologically feasible. However, it is highly likely the species will remain at some risk due to the continued pressure on its habitats from a rapidly growing human population in the Fraser Valley.

<sup>1</sup> Draft Policy on the Feasibility of Recovery, Species at Risk Act Policy. January 2005.

Recovery will involve the establishment and/or maintenance of riffle habitat sufficient to maintain a population in each creek. Some management will be required in all three watersheds. It should focus on in-stream flow protection in Bertrand Creek, restriction of beaver impoundment in Pepin Brook, and the restoration of riffle habitat in Fishtrap Creek. Appropriate recovery actions in the Brunette River are unknown pending population and habitat status surveys.

### **Recovery Goal, Objectives and Corresponding Approaches**

#### ***Recovery Goal:***

To ensure the long-term viability of Nooksack dace populations throughout their natural distribution in Canada.

#### ***Recovery Objectives***

- 1. For all currently and historically suitable habitats in native streams to be occupied by 2015.*

Watershed	Habitat with High Potential Productivity Occupied in 2004 (km)	Total Habitat with High Potential Productivity (km)
Bertrand Creek	<6.5	10.0
Pepin Brook	<2	2.8
Fishtrap Creek	unknown	8.5
Brunette River	unknown	unknown

#### ***Rationale:***

A significant portion of habitat with high potential productivity is not currently occupied, primarily due to riffle degradation or loss to drying, sediment deposition and beaver impoundment. Achievement of interim population recovery targets in the three surveyed watersheds will require that all habitat with high potential productivity be occupied (see objective 2 below). In most cases unoccupied areas could be rendered habitable quickly by increasing water flow, controlling beaver, and/or implementing fish-sensitive drainage maintenance practices.

- 2. To increase Nooksack dace abundance to target levels in all watersheds by 2015.*

Watershed	Area of Riffle in Potential Habitat Reaches (m <sup>2</sup> )	Population Target (excludes young of year)
Bertrand Creek	3000	5700*
Pepin Brook	2300**	4400*
Fishtrap Creek	2030	3900*
Brunette River	unknown	unknown pending habitat survey

\*Assumes an average density of 1.9 Nooksack dace per m<sup>2</sup> riffle in suitable habitat (Inglis et al. 1994). Rounded to nearest hundred.

\*\* Based on 1999 survey. By 2001 approximately 200 m<sup>2</sup> of riffle was lost to beaver ponding (Pearson 2004a).

#### *Rationale*

Ideally population targets would be based on robust population viability analyses. Unfortunately the necessary demographic data is lacking for Nooksack dace. An appropriate guideline for minimum viable population (MVP) size in vertebrate species, based on an extensive review of the scientific literature (Reed et al. 2003, Thomas 1990), is 7000 breeding adults (median value; range 2000-10000). This abundance is considered adequate to maintain genetic diversity and to buffer the population from random variations in survival, and thus to maintain long-term viability in the absence of deterministic factors causing the population to decline.

Populations of Nooksack dace in each of the four watersheds are essentially independent of one another, with extremely low probability of natural exchange of individuals between watersheds because of the very large distances of unsuitable habitat that separate populations. Natural recolonization of habitat from which a population has been extirpated (rescue effect) is therefore highly unlikely. Each watershed, consequently, warrants a separate recovery target in the low to mid thousands.

High quality habitat in Bertrand Creek supported an average of 1.9 dace/m<sup>-2</sup> (n=20, SE = 0.35) in the single available direct estimate of density (Inglis et al. 1994). If all riffle areas in all reaches with habitat with high potential productivity supported this density, total adult abundance would be in the low thousands for each watershed. This suggests that for Nooksack dace in the three surveyed watersheds, the maximum achievable population size is close to the minimum viable population size and that all suitable habitats should be designated critical.

3. To ensure that at least one reach in each watershed supports a high density of Nooksack dace.

#### *Rationale*

Within each watershed, individual populations may be structured as metapopulations, with different subpopulations separated by poor quality habitat, and some level of exchange of individuals between sub-populations. Population persistence in such systems is dependent upon the existence if one or more source areas where population growth is positive and densities are high.

### ***Broad Strategies to be Taken to Address Threats***

Eight broad strategies have been identified in support of the recovery objectives.

1. Protect, create and enhance riffle habitat in habitat reaches with high potential productivity.
2. Establish or maintain adequate baseflow in all habitats with high potential productivity.
3. Reduce sediment entry to creeks.
4. Ensure the integrity and proper functioning of riparian zones throughout watersheds.
5. Reduce habitat fragmentation
6. Encourage stewardship amongst private landowners and the general public.
7. Minimize toxic contamination of creeks.
8. Minimize impacts of introduced predators.

In Table 4 these are prioritized, detailed and related to the relevant recovery goals and objectives.

### ***Evaluation***

Monitoring and evaluation of a subset of populations will occur each year with the status of each population and watershed being evaluated every 5 years at minimum. Performance measures for each objective and broad strategy are listed in Tables 5 and 6. Details and priorities of strategy implementation will be provided in the Action Plan.

### ***Effects on Other Species***

Most recovery efforts will benefit co-occurring native species including steelhead, cutthroat trout, and coho salmon. All three Nooksack dace streams in Canada also contain the Salish sucker (*Catostomus* sp.), which is also listed as Endangered under SARA. Most of the proposed strategies for Nooksack dace recovery should also benefit Salish sucker, although there is potential for conflict over beaver management. In some cases beaver control and dam removal may benefit a Nooksack dace population by restoring riffle habitat, but harm Salish sucker population by eliminating deep pool and marsh habitat. Recovery activities for the two species will be coordinated in watersheds where they co-occur through the development of a multi-species Action Plan. Beaver management will be intended to restore that species' natural balance in these watersheds. Specific measures of controlling beavers and their dams will be determined in the Action Plan.

### ***Approaches to Recovery***

An active adaptive management approach (Walters & Holling 1990) should be used in planning and implementing recovery. Whenever possible management actions should be conducted as controlled experiments designed to inform ongoing recovery and action planning. Recovery planning and implementation should occur at the scale of individual watersheds as their populations are isolated from one another and face differing suites of threats in each watershed.

## **Actions Already Complete or Underway**

### *Landowner Contact and Public Education Programs*

A Recovery Implementation Group (RIG) has been formed. The RIG, in cooperation with local stewardship groups, has developed programs to contact landowners in three Nooksack dace watersheds. A public meeting to exchange information was held in each watershed. In addition, colour display posters on Nooksack dace have been given to stewardship groups in Langley for use in public events.

Table 4: Broad strategies for Nooksack dace recovery and details of associated research and management activities. Underlined points should not be postponed despite lack of full scientific certainty.

Broad Strategy	Obj. No.	Threats Addressed	Priority	Specific Steps	Outcomes or Deliverables
1. Protect, create and enhance riffle habitat in habitat reaches with high potential productivity.	1,2,3	Physical destruction of habitat Riffle loss to beaver ponds. Habitat fragmentation	High	Identify high priority sites for protection, restoration or habitat creation. Assess benefits of riffle creation and enhancement to Nooksack dace populations. Estimate current extent of riffle loss to authorized and unauthorized stream and ditch dredging and to beaver activity. Work with stewardship groups and landowners to identify, and implement habitat creation and restoration projects. Develop best management practices and work plans for habitat reaches with high potential productivity that require drainage maintenance or beaver management.	Protection of habitats with high potential productivity through stewardship agreements, conservation covenants or acquisition of lands containing habitats with high potential productivity Riffle creation/ enhancement projects identified and developed. Public education materials on importance of riffle habitat to fish developed and distributed to landowners. Advice on Nooksack dace habitat requirements and beaver management available to local stewardship groups and agency staff involved in habitat work.
2. Establish and maintain adequate baseflow in all habitats with high potential productivity.	1,2	Seasonal Lack of Water Habitat fragmentation	High	Identify watersheds vulnerable to inadequate baseflow for Nooksack dace. Develop water balance models for watersheds. Establish biologically based minimum in-stream flows for habitats with high potential productivity. Develop wetland restoration projects in vulnerable watersheds. Investigate need and feasibility of supplementing baseflow with well water. Develop and distribute public education materials on impacts of water use on fish and wildlife to landowners and public.	Water balance model showing relative influences of groundwater extraction, surface water extraction, and gravel removal on baseflow for each vulnerable watershed. Objectives for present and future water management in vulnerable watersheds (baseflow and water withdrawal). Adequate water rights for conservation purposes in established and vulnerable watersheds.
3. Reduce	1,2	Sediment	High	Estimate levels of sediment in riffles that	Maximum recommended levels of sediment

sediment entry to creeks.		deposition		is harmful to Nooksack dace. Map, assess and prioritize mitigation for riffle sedimentation in all watersheds. Work with landowners, municipal governments, and stewardship groups to prevent, mitigate and restore sediment degradation of riffles, from urban, agricultural and industrial sources. Develop and distribute public education materials on sediment impacts on fish and wildlife to landowners.	content established for habitat riffles with high potential productivity. Restoration of degraded riffles completed at high priority sites. Mitigation projects to reduce sediment entry completed (e.g. riparian planting, stormsewer retrofits, improved settling ponds).
4. Ensure the integrity and proper functioning of riparian zones throughout watersheds.	1,2,3	Sediment deposition Physical destruction of habitat Toxicity Hypoxia	High	Conduct riparian assessments of habitat reaches with high potential productivity as the basis of proposed riparian buffer widths. Identify, prioritize and develop riparian planting or other projects in cooperation with landowners, stewardship groups and government agencies. Develop and distribute public education materials on riparian reserve strips to landowners	Riparian planting projects completed in high priority areas. Educational materials developed and included in landowner contact programs and other public education applications.
5. Reduce habitat fragmentation.	1,2	Habitat Fragmentation	Med.	Assess the ability of different life history stages to cross different types of barriers. Identify permanent/seasonal barriers and prioritize for mitigation.	Use of strategically located restoration projects to eliminate barriers and provide 'stepping stones' for dispersal to other riffle-rich reaches. Advice on prioritizing restoration projects available to local stewardship groups and agency staff involved in habitat work.
6. Encourage stewardship amongst private landowners, local governments and the general public.			Med.	Give presentations and field tours on Nooksack dace and watershed ecology to local stewardship groups, schools and others. Advise stewardship groups, agency staff, and consultants involved in habitat work on Nooksack dace habitat requirements.	Increased awareness of Nooksack dace and local stream ecology among public.  Nooksack dace habitat features incorporated into in-stream works undertaken for other purposes.

7. Minimize toxic contamination of creeks.	1,2,3	Toxicity	Med.	<p>Estimate extent and severity of toxic contamination of creeks.</p> <p>Work with municipalities to identify, prioritize and develop projects to improve storm water treatment.</p> <p>Increase width and continuity of riparian reserve areas on agricultural lands (see strategy 3).</p> <p>Develop and distribute public education materials on pesticide/herbicide impacts on fish and wildlife to landowners.</p>	<p>Stormwater treatment projects completed at high priority sites.</p> <p>Riparian planting projects completed in high priority areas.</p> <p>Educational materials developed and included in landowner contact programs and other public education applications.</p>
8. Minimize impacts of introduced predators.	1,2,3	Increased predation	Low	<p>Document distribution and densities of introduced predators in each watershed.</p> <p>Assess impact of riffle loss to drying on predation risk.</p> <p>Develop and distribute public education materials on potential impacts of introduced predators on native species to landowners and recreational fishers.</p>	<p>Introduced predator distributions mapped in each watershed.</p> <p>Educational materials developed and included in landowner contact programs and other public education applications.</p>

Table 5: Performance measures for evaluating the achievement of objectives.

<b>Objectives</b>	<b>Process Performance Measure</b>	<b>Biological Performance Measure</b>
1. For all currently and historically utilized habitats in native streams to be occupied by 2015.	Habitat with high potential productivity identified and occupancy evaluated in all watersheds.	Proportion of habitat with high potential productivity occupied.
2. To increase Nooksack dace abundance to target levels in all watersheds by 2015.	Development of a monitoring protocol for population abundance. Abundance surveys completed in all watersheds.	Estimated population size relative to target population.



---

3. To ensure that at least one reach in each watershed supports a high density of Nooksack dace.	Abundance surveys completed in all watersheds.	Number of reaches where catch-per-unit-effort exceeds 0.8 Nooksack dace per standard Gee-trap (24 h set, $n \geq 10$ )
--	--	--

---

Table 6: Performance measures for evaluating the success of broad recovery strategies.

Broad Strategies	Process Performance Measure	Biological Performance Measure
Protect, create and enhance riffle habitat in habitat reaches with high potential productivity.	Area of riffle habitat restored, created or protected. Number of landowners and others reached in public education and consultation programs.	Area of riffle protected, restored or created in habitat reaches with high potential productivity. Establishment or significant growth of populations in habitat reaches with high potential productivity containing protected, created or enhanced riffles.
Establish and maintain adequate baseflow in all habitats with high potential productivity.	Minimum discharges for maintenance of Nooksack dace habitat established in vulnerable watersheds. Discharge monitored in vulnerable watersheds.	Minimum discharges exceeded in vulnerable watersheds.
Reduce sediment entry to creeks.	Major sources of sediment entry to each watershed identified. Major sources of sediment entry addressed.	Area and proportion of habitat with high potential productivity affected by sediment deposition. Establishment or growth of Nooksack dace populations in habitat reaches with high potential productivity where sediment deposition has been addressed.
Ensure the integrity and proper functioning of riparian zones throughout watersheds.	Length and area of riparian habitat restored in each watershed. Proportion of habitat with high potential productivity for which a riparian assessment has been completed. Proportion of habitat with high potential productivity for which the results of a riparian assessment have been adopted.	Length and proportion of habitat with high potential productivity with greater than 5, 10, and 30 m of riparian reserve. Establishment or significant growth of Nooksack dace populations in habitat reaches with high potential productivity with restored riparian reserve strips.
Reduce habitat fragmentation.	Permanent and seasonal barriers to movement mapped in each watershed.	Quantity of habitat reconnected by removal of barriers. Establishment or growth of Nooksack dace populations in habitat reaches with high potential productivity where habitat fragmentation has been addressed.
Encourage stewardship amongst private landowners and the general public.	Number of non-government organizations /individuals involved in recovery activities. Number of stewardship agreements/conservation covenants signed to protect habitat with high potential productivity. Number of landowners and others reached in public education and consultation programs. Length of habitat with high potential productivity protected or restored on private	Establishment or growth of Nooksack dace populations in habitat reaches with high potential productivity on stewarded lands.

---

Minimize toxic contamination of creeks.	land or with public involvement. Sources of toxic contamination identified. Sources of toxic contamination addressed.	Area and proportion of habitat with high potential productivity affected by contamination. Establishment or growth of Nooksack dace populations in habitat reaches with high potential productivity where toxic contamination has been addressed.
Minimize impacts of introduced predators.	Extent of habitat with high potential productivity occupied by introduced predators mapped.	Proportion of habitat with high potential productivity containing introduced predators. Correlation of establishment or growth of Nooksack dace population with introduced predator absence.

---

**Statement of When An Action Plan Will Be Completed.**

A joint Action Plan will be prepared for Nooksack dace and Salish sucker by April 2008. More detailed plans are being prepared for each of the inhabited watersheds as resources and partnership opportunities become available.

## LITERATURE CITED

- Allan, J. D., M. S. Wipfli, J. P. Caouette, A. Prussian, and J. Rodgers. 2003. Influence of streamside vegetation on inputs of terrestrial invertebrates to salmonid food webs. *Canadian Journal of Fisheries and Aquatic Science* **60**:309-320.
- Blueweiss, L., H. Fox, V. Kudzma, D. Nakashima, R. Peters, and S. Sams. 1978. Relationship between body size and some life history parameters. *Oecologia* **37**:257-272.
- Boyle, C. A., L. Lavkulich, H. Schreier, and E. Kiss. 1997. Changes in land cover and subsequent effects on Lower Fraser Basin ecosystems from 1827 to 1990. *Environmental Management* **21**:185-196.
- Brown, G. W., and J. T. Krygier. 1970. Effects of clear-cutting on stream temperature. *Water Resources Research* **6**:1133-1139.
- Burt, A., D. Kramer, K. Nakatsuru, and C. Spry. 1988. The tempo of reproduction in *Hyphessobrycon pulchripinnis* (Characidae) with a discussion on the biology of 'multiple spawning' in fishes. *Environmental Biology of Fishes* **22**:15-27.
- Castelle, A. J., A. W. Johnson, and C. Conolly. 1994. Wetland and stream buffer size requirements - A review. *Journal of Environmental Quality* **23**:878 - 882.
- CCREM. 1987. Canadian water quality guidelines. Canadian Council of Resource and Environment Ministers, Ottawa.
- Corkran, C. C., and C. Thoms 1996. Amphibians of Oregon, Washington, and British Columbia. Lone Pine Publishing, Edmonton.
- Culp, J. M., and R. W. Davies. 1983. An assessment of the effects of streambank clear-cutting on macroinvertebrate communities in a managed watershed. Canadian Technical Reports on Fisheries and Aquatic Science **1208**:115 p.
- Davies, P. E., and M. Nelson. 1994. Relationships between riparian buffer widths and the effects of logging on stream habitat, invertebrate community composition and fish abundance. *Australian Journal of Marine and Freshwater Research* **45**:1289-1305.
- Dhondt, K., P. Boeckx, O. Van Cleemput, G. Hofman, and F. de Troch. 2002. Seasonal groundwater nitrate dynamics in a riparian buffer zone. *Agronomie (Paris)* **22**:747-753.
- Fisheries and Oceans Canada. 1998. Wild, threatened, endangered and lost streams of the lower Fraser Valley Summary Report: Lower Fraser Valley Stream Review Vol. 3. Fraser River Action Plan, Habitat and Enhancement Branch, Fisheries and Oceans Canada, Vancouver.
- Fox, M. G., and A. K. Keast. 1990. Effects of winterkill on population structure and prey consumption patterns of pumpkinseed in isolated beaver ponds. *Canadian Journal of Zoology* **68**:2489-2498.
- Frissell, C. A., W. J. Liss, C. E. Warren, and M. D. Hurley. 1986. A hierarchical framework for stream habitat classification: viewing streams in a watershed context. *Environmental Management* **10**:199-214.
- Gido, K. B., and J. H. Brown. 1999. Invasion of North American drainages by alien fish species. *Freshwater Biology* **42**:387-399.

- Hall, K. J., H. Schreier, and S. J. Brown. 1991. Water quality in the Fraser River basin in J. R. Griggs, editor. *Water is sustainable development: Exploring our common future in the Fraser River Basin*. Westwater Research Centre, University of British Columbia, Vancouver.
- Hanson, W. D., and R. S. Campbell. 1963. The effects of pool size and beaver activity on distribution and abundance of warm-water fishes in a North Missouri stream. *American Midland Naturalist* **69**:136-149.
- Imhof, J. G., J. Fitzgibbon, and W. K. Annable. 1996. A hierarchical evaluation system for characterizing watershed ecosystems for fish habitat. *Canadian Journal of Fisheries and Aquatic Science* **53 (Suppl. 1)**.
- Inglis, S., S. M. Pollard, and M. L. Rosenau. 1994. Distribution and habitat of Nooksack dace (*Rhinichthys sp.*) in Canada. Regional Fisheries Report, B.C. Ministry of Environment, Lands and Parks, Surrey.
- Keast, A. K., and M. G. Fox. 1990. Fish community structure, spatial distribution and feeding ecology in a beaver pond. *Environmental Biology of Fishes* **27**:201-214.
- Kiffney, P. M., J. S. Richardson, and J. P. Bull. 2003. Response of periphyton and insects to experimental manipulation of riparian buffer width along forest streams. *Journal of Applied Ecology* **40**:1060-1076.
- Lavkulich, L. M., K. J. Hall, and H. Schreier. 1999. Land and water interactions: Present and future in M. C. Healey, editor. *Seeking sustainability in the lower Fraser Basin: Issues and Choices*. Institute for Resources and Environment, Westwater Research, University of British Columbia, Vancouver.
- Lee, K. H., T. M. Isenhardt, R. C. Schultz, and S. K. Mickelson. 2003. Multispecies riparian buffers trap sediment and nutrients during rainfall simulations. *Journal of Environmental Quality* **29**:1200-1205.
- Lynch, J. A., G. B. Rishel, and E. S. Corbett. 1984. Thermal alteration of streams draining clearcut watersheds: Quantifications and biological implications. *Hydrobiologia* **111**:161-169.
- Martin, T. L., N. K. Kaushik, J. T. Trevors, and H. R. Whiteley. 1999. Review: denitrification in temperate climate riparian zones. *Water, air and soil pollution* **111**:171-186.
- McKergow, L. A., D. M. Weaver, I. P. Prosser, R. B. Grayson, and A. E. G. Reed. 2003. Before and after riparian management: Sediment and nutrient exports from a small agricultural catchment, Western Australia. *Journal of Hydrology*.
- McPhail, J. D. 1997. Status of the Nooksack dace, *Rhinichthys sp.*, in Canada. *Canadian Field Naturalist* **111**:258-262.
- Miller, R. R., J. D. Williams, and J. E. Williams. 1989. Extinctions of North American fishes during the past century. *Fisheries* **14**:22-38.
- Moring, J. R. 1982. Decrease in stream gravel permeability after clear-cut logging: an indication of intragravel conditions for developing salmonid eggs and alevins. *Hydrobiologia* **88**:295-298.
- Murphy, M. L., J. Heifetz, S. W. Johnson, K. V. Koski, and J. F. Thedinga. 1986. Effects of clear-cut logging with and without buffer strips on juvenile salmonids in Alaskan streams. *Canadian Journal of Fisheries and Aquatic Science* **43**:1521-1533.

- Murphy, M. L., and K. V. Koski. 1989. Input and depletion of woody debris in Alaska streams and implications for streamside management. *North American Journal of Fisheries Management* **9**:427-436.
- National Recovery Team for Salish Sucker and Nooksack Dace. 2005. Critical habitat assessment for Salish sucker and Nooksack dace. Prepared for the British Columbia Ministry of Environment and Fisheries and Oceans Canada by Mike Pearson, Vancouver, B.C.
- Pearson, M. P. 2004a. The ecology, status, and recovery potential of Nooksack dace and Salish sucker in Canada. Ph.D. thesis, University of British Columbia, Vancouver, Canada.
- Pearson, M. P. 2004b. Threats to the Salish sucker and Nooksack dace. Prepared for the National Recovery Team for Salish sucker and Nooksack dace, Fisheries and Oceans Canada, Vancouver.
- Puckett, L. J. 2004. Hydrologic controls on the transport and fate of nitrate in ground water beneath riparian buffer zones: results from thirteen studies across the United States. *Water Science and Technology* **49**:47-53.
- Reed, H. R., J. J. O'Grady, B. W. Brook, J. D. Ballou, and R. Frankham. 2003. Estimates of minimum viable population sizes for vertebrates and factors influencing those estimates. *Biological Conservation* **113**:23-34.
- Richter, B. D. 1997. Threats to imperiled freshwater fauna. *Conservation Biology* **11**:1081-1093.
- Schlosser, I. J. 1991. Stream fish ecology: A landscape perspective. *BioScience* **41**:704-712.
- Schlosser, I. J. 1995. Dispersal, boundary processes and trophic level interactions in streams adjacent to beaver ponds. *Ecology* **76**:908-925.
- Schlosser, I. J., and L. W. Kallemyn. 2000. Spatial variation in fish assemblages across a beaver-influenced successional landscape. *Ecology* **81**:1371-1382.
- Schreier, H., K. J. Hall, L. Elliott, J. Addah, and K. Li. 2003. Ground water and surface water issues in Agassiz, B.C. Institute for Resources, Environment, and Sustainability, University of British Columbia, Vancouver.
- Scott, W. B., and E. J. Crossman 1973. *Freshwater fishes of Canada*. Fisheries Research Board of Canada, Fisheries and Oceans Canada, Ottawa.
- Thomas, C. D. 1990. What do real population dynamics tell us about minimum viable population sizes? *Conservation Biology* **4**:324-327.
- Walters, C. J., and C. S. Holling. 1990. Large-scale management experiments and learning by doing. *Ecology* **71**:2060-2068.
- Waters, T. F. 1995. *Sediment in streams: sources, biological effects and control*. American Fisheries Society Monograph 7, Bethesda, Maryland.
- Weller, D. E., T. E. Jordan, and D. L. Correll. 1998. Heuristic models for material discharge from landscapes with riparian buffers. *Ecological Applications* **8**:1156-1169.
- Whitaker, D. M., A. L. Carroll, and V. A. Montevecchi. 2000. Elevated numbers of flying insects and insectivorous birds in riparian buffer strips. *Canadian Journal of Zoology* **78**:740-747.
- Wigington, P. J., S. M. Griffith, J. A. Field, J. E. Baham, W. R. Horwath, J. Owen, J. H. Davis, S. C. Rain, and J. J. Steiner. 2003. Nitrate removal effectiveness of

a riparian buffer along a small agricultural stream in western Oregon. *Journal of Environmental Quality* **32**:162-170.

**Other Relevant Literature:**

Pearson, M. P. (1998). A review of the distribution, status, and biology of the endangered Salish sucker (*Catostomus sp.*) and Nooksack dace (*Rhinichthys sp.*). Vancouver, B.C. Ministry of Fisheries.

Pearson, M. P. (1998). Habitat inventory and enhancement needs for the endangered Salish sucker (*Catostomus sp.*) and Nooksack dace (*Rhinichthys sp.*). Vancouver, BC Ministry of Fisheries Project Report No. 76.

Pearson, M. P. (2000). The biology and management of Salish sucker and Nooksack dace. The biology and management of species and habitats at risk, Kamloops, B.C., B.C. Ministry of Environment, Lands and Parks, Victoria and University College of the Cariboo, Kamloops.



## **APPENDIX - RECORD OF COOPERATION AND CONSULTATION**

Nooksack dace are listed on Schedule 1 of the Species at Risk Act (SARA), and as an aquatic species are under federal jurisdiction and managed by Fisheries and Oceans Canada (DFO): 200 - 401 Burrard Street, Vancouver, BC.

To assist in the development of an initial draft of this Recovery Strategy, as well as those for other listed freshwater fishes in British Columbia, DFO in cooperation with the Province of BC assembled a group of experts from various levels of government, academia, consultants, and non-governmental organizations to form the Pacific Region Non-Game Freshwater Fish Recovery Team. This team, co-chaired by DFO and the Province of BC, is responsible for drafting recovery strategies for Pacific Region freshwater fish species listed under SARA, including Nooksack dace. In addition, local stakeholders have subsequently established a Recovery Implementation Group for Nooksack dace which has contacted landowners and held public information meetings on the recovery of the species.

Consultation on the draft Recovery Strategy was provided through a series of multi-stakeholder Community Dialogue Sessions and First Nations information exchanges in BC communities, as part of DFO Pacific Region's Fall Consultation Program. A consultation weblink was sent to 198 First Nations, Tribal Councils and Aboriginal Fisheries Commissions, as well as other stakeholders. Notices announcing the Community Dialogue Sessions were placed in 74 newspapers, and announcements specific to Nooksack dace were placed in an additional six newspapers. A specific presentation and discussion session on the proposed Recovery Strategy for Nooksack dace was held in Abbotsford in November 2005, with four attendees. Comments from the session were recorded and archived.

Additional input on the draft Recovery Strategy was sought through a discussion guide and feedback form available on the internet (October – December 2005). No responses were received. Input from the Province of BC and the Township of Langley was received through recovery team participation. An external peer review was requested from several outside experts but no reviews were provided. All feedback received was considered in the finalization of the Recovery Strategy.